

Keep Kansans Healthy & Thriving!

Updated Measles (MMR) Vaccination Recommendations

These updates are for counties that are impacted by the **current outbreak of measles** and adjacent Kansas counties.



INFANTS (6–11 MONTHS)

Get an **early dose of MMR vaccine** (infant dose), then follow CDC's schedule:

- Another dose at 12 through 15 months of age *and*
- A final dose at 4 through 6 years of age

Reminder: The MMRV (measles, mumps, rubella, and **varicella/chickenpox**) vaccine should not be given to children younger than 12 months.



TEENS AND ADULTS

- **Those previously vaccinated with one dose** of MMR vaccine should receive a second dose.
- **Those with no evidence of immunity** should receive one dose of MMR vaccine immediately and follow with a second dose at least 28 days later.



CHILDREN (12+ MONTHS)*

Those who are not yet vaccinated should receive

- One dose immediately *and*
- A second dose at least 28 days after the first

Those previously vaccinated with one dose should receive

- A second dose at least 28 days after the first

* Two doses given after a child's first birthday provides lifelong protection. Children who received an early dose at 6–11 months due to international travel should follow CDC's schedule for a total of three doses:

- Another dose at 12 through 15 months of age (*at least 28 days after first dose*) *and*
- A final dose at 4 through 6 years of age



Questions? Contact your healthcare provider!

IKC IMMUNIZE
KANSAS
COALITION