

"Working together to protect all Kansans from vaccine-preventable diseases."



IMMUNIZE KANSAS COALITION

The Immunize Kansas Coalition is a group of engaged organizations and individuals including healthcare providers, public health professionals, researchers, educators, healthcare payers, associations and nonprofits, vaccine producers, school nurses, pharmacists, parents, grandparents, and more who bring their expertise to support our mission.



staff@immunizekansascoalition.org



@immunizekscoalition



@immunizekscoalition



@ImmKSCoalition



Immunize Kansas Coalition

For information on vaccines, educational resources, and more visit our website!







Fall and winter months typically see an increase in respiratory viruses circulating in the community, which can cause discomfort and, in some cases, serious illness. Vaccination is crucial in preventing flu, RSV, and COVID-19.

Learn more about respiratory viruses, who is most vulnerable, and how vaccines build community immunity!

What is Flu (Influenza)?

Flu is a contagious respiratory virus caused by influenza viruses that infects the nose, throat, and sometimes the lungs.

What is RSV?

RSV usually causes mild, cold-like symptoms. Most people recover in a week or two, but RSV can be serious, especially for babies and older adults.

What is COVID-19?

COVID-19 can be very contagious and spreads quickly. Most people with COVID-19 have mild symptoms, but some people can become severely ill.

What are the Symptoms?

SYMPTOMS	FLU (INFLUENZA)	RSV	COVID-19
Headache or Body Aches	8	×	Θ
Difficulty Breathing	×	\bigcirc	8
Fatigue	8	×	8
Fever	8	Θ	Θ
Loss of Taste or Smell	×	×	Θ
Sore Throat	Θ	×	8
Wheezing	×	8	×

RarelySometimesOften

If you have symptoms of a respiratory virus, there are tests that can help determine which virus you may have and inform your treatment and prevention actions, like physical spacing and mask use.

Source: National
Foundation for Infectious

Diseases (NFID)

Stay Protected with Vaccines!

Flu:

Seasonal flu vaccine is recommended for **everyone over 6 months** (with rare exceptions).



RSV

Protecting Babies:

- RSV vaccine given during pregnancy; if 32–36 weeks pregnant during September–January.
- RSV antibody immunization is recommended for infants under 8
 months born during or entering their first RSV season (if the mother
 was not vaccinated). It's also recommended for some children ages
 8 through 19 months who are at increased risk of RSV and entering
 their second RSV season.

Protecting Older Adults:

 RSV vaccine for adults over 75 years of age and adults ages 60-74 at increased risk of severe RSV.

COVID-19

Those **6 months and older should stay up-to-date** with recommended COVID-19 vaccines.

To learn more about respiratory viruses and how to stay protected, scan the QR code.





Vaccine recommendations may change over time. Please refer to the CDC for the latest recommendations.

scan the QR code.



Talk with your healthcare provider about the vaccines recommended for you.