

Test Your Meningococcal Vaccine Knowledge

1. At any given time, how many people carry the meningococcal bacteria?

- a. 1 in 1000
- b. 1 in 100
- c. 1 in 10
- d. 1 in 2

2. Human beings are the only reservoirs in which meningococcal bacteria can live.

- a. True
- b. False

3. About what percent of confirmed meningococcal disease cases are fatal?

- a. 1%
- b. 2%
- c. 5%
- d. 10%

4. At what age should a person get a meningococcal conjugate vaccination?

- a. 1-2 years old
- b. 11-12 years old
- c. 18-21 years old
- d. 21 years or older

5. What percent of Kansas adolescents currently have at least one of their meningitis vaccinations?

- a. 64%
- b. 72%
- c. 81%
- d. 88%

6. Meningococcal disease is the same as meningitis.

- a. True
- b. False

7. The early symptoms of meningococcal disease often reflect what other illness?

- a. mononucleosis
- b. the flu
- c. foodborne illnesses
- d. chicken pox

8. What is the bacteria that causes meningococcal disease?

- a. *Neisseria meningitidis*
- b. *Escherichia coli*
- c. *Salmonella typhi*
- d. *Listeria monocytogenes*

9. Which of these groups is at a higher risk for contracting meningococcal disease?

- a. Those living in college residence halls
- b. Those traveling to different countries
- c. Those who are going into the military
- d. All of these groups are at higher risk

10. Nationally, ~1 out of 3 cases of meningococcal disease in people 11+ years old are caused by serogroup B.

- a. True
- b. False

11. The following are brand names of meningococcal conjugate vaccines:

- a. M-M-R II and ProQuad
- b. Menactra and Menveo
- c. Prevnar 13 and Pneumovax 23
- d. Gardasil and Cervarix

12. There are several different brands of meningococcal vaccines, but they are all interchangeable.

- a. True
- b. False

Check your answers on the back.

Answers

1. At any given time, how many people carry the meningococcal bacteria?

- a. 1 in 1000
- b. 1 in 100
- c. 1 in 10**
- d. 1 in 2

2. Human beings are the only reservoirs in which meningococcal bacteria can live.

- a. True**
- b. False

3. About what percent of confirmed meningococcal disease cases are fatal?

- a. 1%
- b. 2%
- c. 5%
- d. 10%**

4. At what age should a person get a meningococcal conjugate vaccination?

- a. 1-2 years old
- b. 11-12 years old** – the first vaccination is given at this age, with a booster given at age 16-18
- c. 18-21 years old
- d. 21 years or older

5. What percent of Kansas adolescents currently have at least one of their meningitis vaccinations?

- a. 64%
- b. 72%** - National Immunization Survey-Teen '17
- c. 81%
- d. 88%

6. Meningococcal disease is the same as meningitis.

- a. True
- b. False**

7. The early symptoms of meningococcal disease often reflect what other illness?

- a. mononucleosis
- b. the flu**
- c. foodborne illnesses
- d. chicken pox

8. What is the bacteria that causes meningococcal disease?

- a. *Neisseria meningitidis***
- b. *Escherichia coli*
- c. *Salmonella typhi*
- d. *Listeria monocytogenes*

9. Which of these groups is at a higher risk for contracting meningococcal disease?

- a. Those living in college residence halls
- b. Those traveling to certain countries
- c. Those who are in the military
- d. All of these groups are at higher risk**

10. Nationally, ~1 out of 3 cases of meningococcal disease in people 11+ years old are caused by serogroup B.

- a. True**
- b. False

11. The following are brand names of meningococcal conjugate vaccines:

- a. M-M-R II and ProQuad
- b. Menactra and Menveo**
- c. Prevnar 13 and Pneumovax 23
- d. Gardasil and Cervarix

12. There are several different brands of meningococcal vaccines, but they are all interchangeable.

- a. True
- b. False** – There are multiple brands, but they are not interchangeable. Talk with your health care provider to ensure you remain on the correct series.